

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 1 174 508 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
23.01.2002 Bulletin 2002/04

(51) Int Cl.7: **C12N 15/52, C12N 15/54,  
C12P 13/14, C12R 1/13,  
C12R 1/15**

(21) Application number: **01115635.3**

(22) Date of filing: **03.07.2001**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR**  
Designated Extension States:  
**AL LT LV MK RO SI**

- Nakamura, Jun, Ajinomoto Co., Inc.  
Kawasaki-ku, Kawasaki-shi, Kanagawa (JP)
- Izui, Hiroshi, Ajinomoto Co., Inc.  
Kawasaki-ku, Kawasaki-shi, Kanagawa (JP)
- Nakamatsu, Tsuyoshi, Ajinomoto Co., Inc.  
Kawasaki-ku, Kawasaki-shi, Kanagawa (JP)

(30) Priority: **05.07.2000 JP 2000204256**

(71) Applicant: **Ajinomoto Co., Inc.**  
**Tokyo 104 (JP)**

(74) Representative: **HOFFMANN - EITLE**  
**Patent- und Rechtsanwälte Arabellastrasse 4**  
**81925 München (DE)**

(72) Inventors:  
• Ohtaki, Hiromi, Ajinomoto Co., Inc.  
Kawasaki-ku, Kawasaki-shi, Kanagawa (JP)

(54) **Bacterium producing L-glutamic acid and method for producing L-glutamic acid**

(57) L-Glutamic acid is produced by culturing a co-ryneform bacterium having L-glutamic acid producing ability, in which trehalose synthesis ability is decreased or deleted by, for example, disrupting a gene coding for

trehalose-6-phosphate synthase, a gene coding for maltotigosyltrehalose synthase, or both of these genes to produce and accumulate L-glutamic acid in the medium, and collecting the L-glutamic acid from the medium.

**EP 1 174 508 A2**

**Description**Background of the invention5 Field of the Invention

[0001] The present invention relates to a novel L-glutamic acid producing bacterium and a method for producing L-glutamic acid by fermentation utilizing it. L-glutamic acid is an important amino acid as foodstuffs, drugs and so forth.

10 Description of the Related Art

[0002] Conventionally, L-glutamic acid is mainly produced by fermentative methods using so-called L-glutamic acid producing coryneform bacteria belonging to the genus *Brevibacterium*, *Corynebacterium* or *Microbacterium*, or mutant strains thereof (Amino Acid Fermentation, pp.195-215, Gakkai Shuppan Center, 1986).

15 [0003] It is known that, in the production of L-glutamic acid by fermentation, trehalose is also produced as a secondary product. Therefore, techniques have been developed for decomposing or metabolizing the produced trehalose. Such techniques include the method of producing an amino acid by fermentation using a coryneform bacterium in which proliferation ability on trehalose is induced (Japanese Patent Laid-open (Kokai) No. 5-276935) and the method of producing amino acid by fermentation using a coryneform bacterium in which a gene coding for trehalose catabolic enzyme is amplified (Korean Patent Publication (B1) No. 165836). However, it is not known how to suppress the formation of trehalose itself in an L-glutamic acid producing bacterium.

[0004] In *Escherichia coli*, the synthesis of trehalose is catalyzed by trehalose-6-phosphate synthase. This enzyme is known to be encoded by *otsA* gene. Further, it has been also reported that an *otsA* gene-disrupted strain of *Escherichia coli* can scarcely grow in a hyperosmotic medium (H.M. Glaever, *et al.*, *J. Bacteriol.*, 170(6), 2841-2849 (1998)). However, the relationship between disruption of *otsA* gene and production of substances has not been known.

25 [0005] On the other hand, although the *treY* gene is known for *Brevibacterium helvolum* among bacteria belonging to the genus *Brevibacterium* bacteria, any *otsA* gene is not known for them. As for bacteria belonging to the genus *Mycobacterium* bacteria, there is known a pathway via a reaction catalyzed by a product encoded by *treS* gene (trehalose synthase (TreS)), which gene is different from the *otsA* gene and *treY* gene, as a gene coding for an enzyme in trehalose biosynthesis pathway (De Smet K.A., *et al.*, *Microbiology*, 146 (1), 199-208 (2000)). However, this pathway utilizes maltose as a substrate and does not relate to usual L-glutamic acid fermentation that utilizes glucose, fructose or sucrose as a starting material.

30 SUMMARY OF THE INVENTION

35 [0006] An object of the present invention is to improve production efficiency of L-glutamic acid in L-glutamic acid production by fermentation using coryneform bacteria through suppression of the production of trehalose as a secondary product.

[0007] The inventors of the present invention assiduously studied in order to achieve the aforementioned object. As a result, they found that bacterium belonging to the genus *Brevibacterium* contained *otsA* gene and *treY* gene like *Mycobacterium tuberculosis* and the production efficiency of L-glutamic acid was improved by disrupting at least one of these genes. Thus, they accomplished the present invention.

[0008] That is, the present invention provides the followings.

45 (1) A coryneform bacterium having L-glutamic acid producing ability, wherein trehalose synthesis ability is decreased or deleted in the bacterium.

(2) The coryneform bacteria according to (1), wherein the trehalose synthesis ability is decreased or deleted by introducing a mutation into a chromosomal gene coding for an enzyme in a trehalose synthesis pathway or disrupting the gene.

50 (3) The coryneform bacteria according to (2), wherein the gene coding for the enzyme in trehalose synthesis pathway consists of a gene coding for trehalose-6-phosphate synthase, a gene coding for maltotriose synthase, or both of these genes.

(4) The coryneform bacteria according to (3), wherein the gene coding for trehalose-6-phosphate synthase codes for the amino acid sequence of SEQ ID NO: 30, and the gene coding for maltotriose synthase codes for the amino acid sequence of SEQ ID NO: 32.

55 (5) A method for producing L-glutamic acid comprising culturing a coryneform bacterium according to any one of (1) to (4) in a medium to produce and accumulate L-glutamic acid in the medium, and collecting the L-glutamic acid from the medium.

(6) A DNA coding for a protein defined in the following (A) or (B):

- (A) a protein having the amino acid sequence of SEQ ID NO: 30,
- (B) a protein having an amino acid sequence of SEQ ID NO: 30 including substitution, deletion, insertion or addition of one or several amino acid residues and having trehalose-6-phosphate synthase activity.

(7) A DNA according to (6), which is a DNA defined in the following (a) or (b):

- (a) a DNA containing a nucleotide sequence comprising at least the residues of nucleotide numbers 484-1938 in the nucleotide sequence of SEQ ID NO: 29,
- (b) a DNA hybridizable with a nucleotide sequence comprising at least the residues of nucleotide numbers 484-1938 in the nucleotide sequence of SEQ ID NO: 29 under a stringent condition, showing homology of 55% or more to the foregoing nucleotide sequence, and coding for a protein having trehalose-6-phosphate synthase activity.

(8) A DNA coding for a protein defined in the following (A) or (B):

- (A) a protein having the amino acid sequence of SEQ ID NO: 32,
- (B) a protein having an amino acid sequence of SEQ ID NO: 32 including substitution, deletion, insertion or addition of one or several amino acid residues and having maltotriosyltrehalose synthase activity.

(9) A DNA according to (8), which is a DNA defined in the following (a) or (b):

- (a) a DNA containing a nucleotide sequence comprising at least the residues of nucleotide numbers 82-2514 in the nucleotide sequence of SEQ ID NO: 31,
- (b) a DNA hybridizable with a nucleotide sequence comprising at least the residues of nucleotide numbers 82-2514 in the nucleotide sequence of SEQ ID NO: 31 under a stringent condition, showing homology of 60% or more to the foregoing nucleotide sequence, and coding for a protein having maltotriosyltrehalose synthase activity.

[0009] The trehalose-6-phosphate synthase activity means an activity to catalyze a reaction in which  $\alpha,\alpha$ -trehalose-6-phosphate and UDP are produced from UDP-glucose and glucose-6-phosphate, and the maltotriosyltrehalose synthase activity means an activity to catalyze a reaction in which maltotriosyltrehalose is produced from maltopentose.

[0010] According to the present invention, production efficiency of L-glutamic acid in L-glutamic acid production by fermentation using coryneform bacteria can be improved through inhibition of the production of trehalose as a secondary product.

#### Preferred Embodiments of the Invention

[0011] Hereafter, the present invention will be explained in detail.

[0012] The coryneform bacterium of the present invention is a coryneform bacterium having L-glutamic acid producing ability, in which trehalose synthesis ability is decreased or deleted.

[0013] The coryneform bacteria referred to in the present invention include the group of microorganisms defined in Bergey's Manual of Determinative Bacteriology, 8th edition, p.599 (1974), which are aerobic Gram-positive rods having no acid resistance and no spore-forming ability aerobic. They have hitherto been classified into the genus *Brevibacterium*, but united into the genus *Corynebacterium* at present (*Int. J. Syst. Bacteriol.*, 41, 255 (1981)), and include bacteria belonging to the genus *Brevibacterium* or *Microbacterium* closely relative to the genus *Corynebacterium*. Examples of such coryneform bacteria are mentioned below.

*Corynebacterium acetoacidophilum*  
*Corynebacterium acetoglutamicum*  
*Corynebacterium alkanolyticum*  
*Corynebacterium callunae*  
*Corynebacterium glutamicum*  
*Corynebacterium lilium* (*Corynebacterium glutamicum*)  
*Corynebacterium melassecola*  
*Corynebacterium thermoaminogenes*  
*Corynebacterium herculis*

*Brevibacterium divaricatum* (*Corynebacterium glutamicum*)  
*Brevibacterium flavum* (*Corynebacterium glutamicum*)  
*Brevibacterium immariophilum*  
*Brevibacterium lactofermentum* (*Corynebacterium glutamicum*)  
5 *Brevibacterium roseum*  
*Brevibacterium saccharolyticum*  
*Brevibacterium thiogenitalis*  
*Brevibacterium ammoniagenes* (*Corynebacterium ammoniagenes*)  
*Brevibacterium album*  
10 *Brevibacterium cerium*  
*Microbacterium ammoniaphilum*

[0014] Specifically, the following strains can be exemplified.

15 *Corynebacterium acetoacidophilum* ATCC 13870  
*Corynebacterium acetoglutamicum* ATCC 15806  
*Corynebacterium alkanolyticum* ATCC21511  
*Corynebacterium callunae* ATCC 15991  
*Corynebacterium glutamicum* ATCC 13020, 13032, 13060  
20 *Corynebacterium lilium* (*Corynebacterium glutamicum*) ATCC 15990  
*Corynebacterium melassecola* ATCC 17965  
*Corynebacterium thermoaminogenes* AJ12340 (FERM BP-1539)  
*Corynebacterium herculis* ATCC13868  
*Brevibacterium divaricatum* (*Corynebacterium glutamicum*) ATCC 14020  
25 *Brevibacterium flavum* (*Corynebacterium glutamicum*) ATCC 13826, ATCC 14067  
*Brevibacterium immariophilum* ATCC 14068  
*Brevibacterium lactofermentum* (*Corynebacterium glutamicum*) ATCC 13665, ATCC 13869  
*Brevibacterium roseum* ATCC 13825  
*Brevibacterium saccharolyticum* ATCC 14066  
30 *Brevibacterium thiogenitalis* ATCC 19240  
*Brevibacterium ammoniagenes* (*Corynebacterium ammoniagenes*) ATCC 6871  
*Brevibacterium album* ATCC 15111  
*Brevibacterium cerium* ATCC 15112  
*Microbacterium ammoniaphilum* ATCC 15354  
35

[0015] The trehalose synthesis ability of such coryneform bacteria as mentioned above can be decreased or deleted by mutagenizing or disrupting a gene coding for an enzyme in trehalose synthesis pathway using mutagenesis treatment or genetic recombination technique. Such a mutation may be a mutation that suppresses transcription or translation of the gene coding for the enzyme in trehalose synthesis pathway, or a mutation that causes elimination or decrease of an enzyme in trehalose synthesis pathway. The enzyme in trehalose synthesis pathway may be exemplified by, for example, trehalose 6-phosphate synthase, maltoseglycotrehalose synthase, or both of these.

[0016] The disruption of a gene coding for an enzyme in trehalose synthesis pathway can be performed by gene substitution utilizing homologous recombination. A gene on a chromosome of a coryneform bacterium can be disrupted by transforming the coryneform bacterium with DNA containing a gene coding for an enzyme in trehalose synthesis pathway modified so that a part thereof should be deleted and hence the enzyme in trehalose synthesis pathway should not normally function (deletion type gene), and allowing recombination between the deletion type gene and a normal gene on the chromosome. Such gene disruption by homologous recombination has already been established. To this end, there can be mentioned a method utilizing a linear DNA or a cyclic DNA that does not replicate in coryneform bacteria and a method utilizing a plasmid containing a temperature sensitive replication origin. However, a method utilizing a cyclic DNA that does not replicate in coryneform bacteria or a plasmid containing a temperature sensitive replication origin is preferred.

[0017] The gene coding for an enzyme in trehalose synthesis pathway may be exemplified by, for example, the *otsA* gene or *treY* gene, or it may consist of both of these. Since the nucleotide sequences of the *otsA* gene and *treY* gene of *Brevibacterium lactofermentum* and flanking regions thereof have been elucidated by the present invention, those genes can be easily obtained by preparing primers based on the sequences and performing PCR (polymerase chain reaction, see White, T.J. *et al.*, *Trends Genet.*, 5, 185 (1989)) using the primers and chromosomal DNA of *Brevibacterium lactofermentum* as a template.

[0018] The nucleotide sequence comprising the *otsA* gene and the nucleotide sequence comprising the *treY* gene

of *Brevibacterium lactofermentum* obtained in the examples described later are shown in SEQ ID NOS: 29 and 31, respectively. Further, the amino acid sequences encoded by these nucleotide sequences are shown in SEQ ID NOS: 30 and 32, respectively.

[0019] The *otsA* gene and *treY* gene each may be one coding for a protein including substitution, deletion, insertion or addition of one or several amino acids at one or a plurality of positions, provided that the activity of trehalose-6-phosphate synthase or maltotoligosyltrehalose synthase encoded thereby is not deteriorated. While the number of "several" amino acids differs depending on positions or types of amino acid residues in the three-dimensional structure of the protein, it is preferably 1-40, more preferably 1-20, further preferably 1-10.

[0020] A DNA coding for the substantially same protein as trehalose-6-phosphate synthase or maltotoligosyltrehalose synthase described above can be obtained by, for example, modifying each of the nucleotide sequences by, for example, the site-directed mutagenesis method so that one or more amino acid residues at a specified site should involve substitution, deletion, insertion, addition or inversion. Such a DNA modified as described above may also be obtained by a conventionally known mutation treatment. The mutation treatment includes a method of treating DNA coding for trehalose-6-phosphate synthase or maltotoligosyltrehalose *in vitro*, for example, with hydroxylamine, and a method for treating a microorganism, for example, a bacterium belonging to the genus *Escherichia* harboring a DNA coding for trehalose-6-phosphate synthase or maltotoligosyltrehalose with ultraviolet irradiation or a mutating agent usually used for mutation treatment such as N-methyl-N'-nitro-N-nitrosoguanidine (NTG) and nitrous acid.

[0021] The substitution, deletion, insertion, addition, or inversion of nucleotide as described above also includes a naturally occurring mutant or variant on the basis of, for example, individual difference or difference in species or genus of microorganisms that harbor trehalose-6-phosphate synthase or maltotoligosyltrehalose.

[0022] A DNA coding for the substantially same protein as trehalose-6-phosphate synthase or maltotoligosyltrehalose synthase described above can be obtained by expressing such a DNA having a mutation as described above in a suitable cell, and examining the trehalose-6-phosphate synthase activity or maltotoligosyltrehalose synthase activity of the expression product.

[0023] A DNA coding for substantially the same protein as trehalose-6-phosphate synthase can also be obtained by isolating a DNA hybridizable with a DNA having, for example, a nucleotide sequence corresponding to nucleotide numbers of 484-1938 of the nucleotide sequence shown in SEQ ID NO: 29 or a probe that can be prepared from the nucleotide sequence under a stringent condition, showing homology of 55% or more, preferably 65% or more, more preferably 75% or more, to the foregoing nucleotide sequence, and having trehalose-6-phosphate synthase activity from a DNA coding for trehalose-6-phosphate synthase having a mutation or from a cell harboring it. Similarly, a DNA coding for substantially the same protein as maltotoligosyltrehalose synthase can also be obtained by isolating a DNA hybridizable with a DNA having, for example, a nucleotide sequence corresponding to nucleotide numbers of 82-2514 of the nucleotide sequence shown in SEQ ID NO: 31 or a probe that can be prepared from the nucleotide sequence under a stringent condition, showing homology of 60% or more, preferably 70% or more, more preferably 80% or more, to the foregoing nucleotide sequence, and having maltotoligosyltrehalose synthase activity from a DNA coding for maltotoligosyltrehalose synthase having a mutation or from a cell harboring it.

[0024] The "stringent condition" referred to herein is a condition under which so-called specific hybrid is formed, and non-specific hybrid is not formed. It is difficult to clearly express this condition by using any numerical value. However, for example, the stringent condition includes a condition under which DNA's having high homology, for example, DNA's having homology of not less than 55%, preferably not less than 60%, are hybridized with each other, and DNA's having homology lower than the above level are not hybridized with each other. Alternatively, the stringent condition is exemplified by a condition under which DNA's are hybridized with each other at a salt concentration corresponding to an ordinary condition of washing in Southern hybridization, i.e., 1 x SSC, 0.1 % SDS, preferably 0.1 x SSC, 0.1 % SDS, at 60°C.

[0025] As the probe, a partial sequence of each gene can also be used. Such a probe can be produced by PCR using oligonucleotides produced based on the nucleotide sequence of each gene as primers and a DNA fragment containing each gene as a template. When a DNA fragment in a length of about 300 bp is used as the probe, the washing conditions for the hybridization may consists of 50°C, 2 x SSC and 0.1% SDS.

[0026] Genes hybridizable under such conditions as described above include those having a stop codon generated in a coding region of the genes, and those having no activity due to mutation of active center. However, such mutants can be easily removed by ligating each of the genes with a commercially available expression vector, and measuring trehalose-6-phosphate synthase activity or maltotoligosyltrehalose synthase activity.

[0027] When an *otsA* gene or *treY* gene is used for the disruption of these genes on chromosomes of coryneform bacteria, the encoded trehalose-6-phosphate synthase or maltotoligosyltrehalose synthase are not required to have their activities. Further, the *otsA* gene or *treY* gene used for the gene disruption may be a gene derived from another microorganism, so long as they can undergo homologous recombination with these genes of coryneform bacteria. For example, an *otsA* gene of bacterium belonging to the genus *Escherichia* or *Mycobacterium*, *treY* gene of bacterium belonging to the genus *Arthrobacter*, *Brevibacterium helvolum*, or bacterium belonging to the genus *Rhizobium* can

be mentioned.

[0028] A deletion type gene of the *otsA* gene or *treY* gene can be prepared by excising a certain region with restriction enzyme(s) from a DNA fragment containing one of these genes or a part of them to delete at least a part of coding region or an expression regulatory sequence such as promoter.

[0029] Further, a deletion type gene can also be obtained by performing PCR using primers designed so that a part of gene should be deleted. Furthermore, a deletion type gene may be one obtained by single nucleotide mutation, for example, a frame shift mutation.

[0030] Gene disruption of the *otsA* gene will be explained hereafter. Gene disruption of the *treY* gene can be performed similarly.

[0031] An *otsA* gene on a host chromosome can be replaced with a deletion type *otsA* gene as follows. That is, a deletion type *otsA* gene and a marker gene for resistance to a drug, such as kanamycin, chloramphenicol, tetracycline and streptomycin, are inserted into a plasmid that cannot autonomously replicate in coryneform bacteria to prepare a recombinant DNA. A coryneform bacterium can be transformed with the recombinant DNA, and the transformant strain can be cultured in a medium containing the drug to obtain a transformant strain in which the recombinant DNA was introduced into chromosomal DNA. Alternatively, such a transformant strain can be obtained by using a temperature sensitive plasmid as the plasmid, and culturing the transformants at a temperature at which the temperature sensitive plasmid cannot replicate.

[0032] In a strain in which the recombinant DNA is incorporated into a chromosome as described above, the recombinant DNA causes recombination with an *otsA* gene sequence that originally exists on the chromosome, and two of fused genes comprising the chromosomal *otsA* gene and the deletion type *otsA* gene are inserted into the chromosome so that other portions of the recombinant DNA (vector portion and drug resistance marker gene) should be interposed between them.

[0033] Then, in order to leave only the deletion type *otsA* gene on the chromosomal DNA, one copy of the *otsA* gene is eliminated from the chromosomal DNA together with the vector portion (including the drug resistance marker gene) by recombination of two of the *otsA* genes. In that case, the normal *otsA* gene is left on the chromosomal DNA and the deletion type *otsA* gene is excised, or conversely, the deletion type *otsA* gene is left on the chromosomal DNA and the normal *otsA* gene is excised. It can be confirmed which type of the gene is left on the chromosomal DNA by investigating structure of the *otsA* gene on the chromosome by PCR, hybridization or the like.

[0034] The coryneform bacterium used for the present invention may have enhanced activity of an enzyme that catalyzes the biosynthesis of L-glutamic acid in addition to the deletion or decrease of trehalose synthesis ability. Examples of the enzyme that catalyzes the biosynthesis of L-glutamic acid include glutamate dehydrogenase, glutamine synthetase, glutamate synthase, isocitrate dehydrogenase, aconitate hydratase, citrate synthase, pyruvate carboxylase, phosphoenolpyruvate carboxylase, phosphoenolpyruvate synthase, enolase, phosphoglyceromutase, phosphoglycerate kinase, glyceraldehyde-3-phosphate dehydrogenase, triosephosphate isomerase, fructose biphosphate aldolase, phosphofructokinase, glucose phosphate isomerase and so forth.

[0035] Further, in the coryneform bacterium used for the present invention, an enzyme that catalyzes a reaction for generating a compound other than L-glutamic acid by branching off from the biosynthetic pathway of L-glutamic acid may be declined or made deficient. Examples of such an enzyme include  $\alpha$ -ketoglutarate dehydrogenase, isocitrate lyase, phosphate acetyltransferase, acetate kinase, acetohydroximate synthase, acetolactate synthase, formate acetyltransferase, lactate dehydrogenase, L-glutamate decarboxylase, 1-pyrroline dehydrogenase and so forth.

[0036] Furthermore, by introducing a temperature sensitive mutation for a biotin activity inhibiting substance such as surface active agents into a coryneform bacterium having L-glutamic acid producing ability, the bacterium becomes to be able to produce L-glutamic acid in a medium containing an excessive amount of biotin in the absence of a biotin activity inhibiting substance (see WO96/06180). As such a coryneform bacterium, the *Brevibacterium lactofermentum* AJ13029 strain disclosed in WO96/06180 can be mentioned. The AJ13029 strain was deposited at the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology (currently, the independent administrative corporation, National Institute of Advanced Industrial Science and Technology, International Patent Organism Depository (Chuo Dai-6, 1-1 Higashi 1-Chome, Tsukuba-shi, Ibaraki-ken, Japan, postal code: 305-5466) on September 2, 1994, and received an accession number of FERM P-14501. Then, it was transferred to an international deposit under the provisions of the Budapest Treaty on August 1, 1995, and received an accession number of FERM BP-5189.

[0037] When a coryneform bacterium having L-glutamic acid producing ability, in which trehalose synthesis ability is decreased or deleted, is cultured in a suitable medium, L-glutamic acid is accumulated in the medium.

[0038] The medium used for producing L-glutamic acid is a usual medium that contains a carbon source, a nitrogen source, inorganic ions and other organic trace nutrients as required. As the carbon source, it is possible to use sugars such as glucose, lactose, galactose, fructose, sucrose, maltose, blackstrap molasses and starch hydrolysate; alcohols such as ethanol and inositol; or organic acids such as acetic acid, fumaric acid, citric acid and succinic acid.

[0039] As the nitrogen source, there can be used inorganic ammonium salts such as ammonium sulfate, ammonium

nitrate, ammonium chloride, ammonium phosphate and ammonium acetate, ammonia, organic nitrogen such as peptone, meat extract, yeast extract, corn steep liquor and soybean hydrolysate, ammonia gas, aqueous ammonia and so forth.

[0040] As the inorganic ions (or sources thereof), added is a small amount of potassium phosphate, magnesium sulfate, iron ions, manganese ions and so forth. As for the organic trace nutrients, it is desirable to add required substances such as vitamin B<sub>1</sub>, yeast extract and so forth in a suitable amount as required.

[0041] The culture is preferably performed under an aerobic condition performed by shaking, stirring for aeration or the like for 16 to 72 hours. The culture temperature is controlled to be at 30°C to 45°C, and pH is controlled to be 5 to 9 during the culture. For such adjustment of pH, inorganic or organic acidic or alkaline substances, ammonia gas and so forth can be used.

[0042] Collection of L-glutamic acid from fermentation broth can be performed by, for example, methods utilizing ion exchange resins, crystallization and so forth. Specifically, L-glutamic acid can be adsorbed on an anion exchange resin and isolated from it, or crystallized by neutralization.

## EXAMPLES

[0043] Hereafter, the present invention will be explained more specifically with reference to the following examples.

### Example 1: Construction of *otsA* gene-disrupted strain of *Brevibacterium lactofermentum*

#### <1> Cloning of *otsA* gene

[0044] Since *otsA* gene of *Brevibacterium lactofermentum* was not known, it was obtained by utilizing a nucleotide sequence of *otsA* gene of another microorganism for reference. The *otsA* genes of *Escherichia* and *Mycobacterium* had been hitherto elucidated for their entire nucleotide sequences (Kaasen I., *et al.*, *Gene*, **145** (1), 9-15 (1994); De Smet K.A., *et al.*, *Microbiology*, **146** (1), 199-208 (2000)). Therefore, referring to an amino acid sequence deduced from these nucleotide sequences, DNA primers P1 (SEQ ID NO: 1) and P2 (SEQ ID NO: 2) for PCR were synthesized first. The DNA primers P1 and P2 corresponded to the regions of the nucleotide numbers of 1894-1913 and 2531-2549 of the nucleotide sequence of the *otsA* gene of *Escherichia coli* (GenBank accession X69160), respectively. They also corresponded to the regions of the nucleotide numbers 40499-40518 and 41166-41184 of the *otsA* gene of *Mycobacterium tuberculosis* (GenBank accession Z95390), respectively.

[0045] Then, PCR was performed by using the primers P1 and P2 and chromosomal DNA of *Brevibacterium lactofermentum* ATCC 13869 as a template with a cycle consisting of reactions at 94°C for 0.5 minute, 50°C for 0.5 minute and 72°C for 4 minutes, which was repeated for 30 cycles. As a result, a substantially single kind of amplified fragment of about 0.6 kbp was obtained. This amplified fragment was cloned into a plasmid vector pCR2.1 by using "Original TA Cloning Kit" produced by Invitrogen to obtain pCotsA. Then, the nucleotide sequence of the cloned fragment was determined.

[0046] Based on the nucleotide sequence of the partial fragment of *otsA* gene obtained as described above, DNA primers P10 (SEQ ID NO: 8) and P12 (SEQ ID NO: 10) were newly synthesized, and unknown regions flanking to the partial fragment was amplified by "inverse PCR" (Triglia, T. *et al.*, *Nucleic Acids Res.*, **16**, 81-86 (1988); Ochman H., *et al.*, *Genetics*, **129**, 921-926 (1990)). The chromosomal DNA of *Brevibacterium lactofermentum* ATCC 13869 was digested with a restriction enzyme *Bam*HI, *Bgl*II, *Clal*, *Hind*III, *Kpn*I, *Mlu*I, *Mun*I, *Sal*I or *Xho*I, and self-ligated by using T4 DNA ligase (Takara Shuzo). By using resultant DNA as a template and the DNA primers P10 and P12, PCR was performed with a cycle consisting of reactions at 94°C for 0.5 minute, 55°C for 1 minute and 72°C for 4 minutes, which was repeated for 30 cycles. As a result, when *Clal* or *Bgl*II was used as the restriction enzyme, an amplified fragment of 4 kbp was obtained for each case. The nucleotide sequences of these amplified fragments were directly determined by using the DNA primers P5 to P9 (SEQ ID NOS: 3-7) and P11 to P15 (SEQ ID NOS: 9-13). Thus, the entire nucleotide sequence of *otsA* gene of *Brevibacterium lactofermentum* ATCC 13869 was determined as shown in SEQ ID NO: 29. The amino acid sequence encoded by this nucleotide sequence is shown in SEQ ID NOS: 29 and 30.

[0047] When homology of the sequence of the aforementioned *otsA* gene was determined with respect to the *otsA* gene of *Escherichia coli* (GenBank accession X69160) and the *otsA* gene of *Mycobacterium tuberculosis* (GenBank accession Z95390), the nucleotide sequence showed homologies of 46.3% and 55.9%, respectively, and the amino acid sequence showed homologies of 30.9% and 51.7%, respectively. The homologies were calculated by using software, "GENETIX-WIN" (Software Development), based on the Lipman-Person method (*Science*, **227**, 1435-1441 (1985)).



<2> Preparation of plasmid for *otsA* gene disruption

[0048] In order to examine presence or absence of improvement effect in L-glutamic acid productivity by disruption of a gene coding for an enzyme in trehalose biosynthesis pathway in coryneform bacteria, a plasmid for *otsA* gene disruption was produced. A plasmid for *otsA* gene disruption was produced as follows. PCR was performed by using the plasmid pCotsA previously constructed in the cloning of the *otsA* gene as a template and the primers P29 (SEQ ID NO: 33) and P30 (SEQ ID NO: 34) comprising *Cla*I site with a cycle consisting of reactions at 94°C for 0.5 minute, 55°C for 0.5 minute and 72°C for 8 minutes, which was repeated for 30 cycles. The amplified fragment was digested with *Cla*I, blunt-ended by using T4 DNA polymerase (Takara Shuzo), and self-ligated by using T4 ligase (Takara Shuzo) to construct a plasmid pCotsAC containing the *otsA* gene having a frame shift mutation (1258-1300th nucleotides of SEQ ID NO: 29 were deleted) at an approximately central part thereof.

<3> Preparation of *otsA* gene-disrupted strain

[0049] By using the plasmid pCotsAC for gene disruption, a L-glutamic acid producing bacterium, *Brevibacterium lactofermentum* ATCC 13869, was transformed by the electric pulse method, and transformants were selected as to the ability to grow in CM2B medium containing 20 mg/L of kanamycin. Because the plasmid pCotsAC for *otsA* gene disruption did not have a replication origin that could function in *Brevibacterium lactofermentum*, resultant transformants obtained by using the plasmid suffered homologous recombination occurred between the *otsA* genes on the chromosome of *Brevibacterium lactofermentum* and the plasmid pCotsAC for gene disruption. From the homologous recombinant strains obtained as described above, strains in which the vector portion of the plasmid pCotsAC for gene disruption was eliminated due to re-occurrence of homologous recombination were selected based on acquired kanamycin sensitivity as a marker.

[0050] From the strains obtained as described above, a strain introduced with the desired frame shift mutation was selected. Selection of such a strain was performed by PCR using chromosomal DNA extracted from a strain that became kanamycin sensitive as a template and the DNA primers P8 (SEQ ID NO: 14) and P13 (SEQ ID NO: 11) with a cycle consisting of reactions at 94°C for 0.5 minute, 55°C for 0.5 minute and 72°C for 1 minutes, which was repeated for 30 cycles, and sequencing of the obtained amplified fragment using the DNA primer P8 to confirm disfunction of the *otsA* gene due to introduction of frame shift mutation. The strain obtained as described above was designated as AOA strain.

Example 2: Construction of *treY* gene-disrupted strain<1> Cloning of *treY* gene

[0051] Since *treY* gene of *Brevibacterium lactofermentum* was not known, it was obtained by using nucleotide sequences of *treY* genes of the other microorganisms for reference. The nucleotide sequences of *treY* genes were hitherto elucidated for the genera *Arthrobacter*, *Brevibacterium* and *Rhizobium* (Maruta K., et al., *Biochim. Biophys. Acta*, 1289 (1), 10-13 (1996); Genbank accession AF039919; Maruta K., et al., *Biosci. Biotechnol. Biochem.*, 60 (4), 717-720 (1996)). Therefore, referring to an amino acid sequence deduced from these nucleotide sequences, the PCR DNA primers P3 (SEQ ID NO: 14) and P4 (SEQ ID NO: 15) were synthesized first. The DNA primers P3 and P4 correspond to the regions of the nucleotide numbers of 975-992 and 2565-2584 of the nucleotide sequence of the *treY* gene of *Arthrobacter* species (GenBank accession D63343), respectively. Further, they correspond to the regions of the nucleotide numbers 893-910 and 2486-2505 of the *treY* gene of *Brevibacterium helvolum* (GenBank accession AF039919), respectively. Furthermore, they correspond to the regions of the nucleotide numbers of 862-879 and 2452-2471 of *treY* gene of *Rhizobium* species (GenBank accession D78001).

[0052] Then, PCR was performed by using the primers P3 and P4 and chromosomal DNA of *Brevibacterium lactofermentum* ATCC13869 as a template with a cycle consisting of reactions at 94°C for 0.5 minute, 55°C for 0.5 minute and 72°C for 2 minutes, which was repeated for 30 cycles. As a result, a substantially single kind of an amplified fragment of about 1.6 kbp was obtained. This amplified fragment was cloned into a plasmid vector pCR2.1 by using "Original TA Cloning Kit" produced by Invitrogen. Then, the nucleotide sequence was determined for about 0.6 kb.

[0053] Based on the nucleotide sequence of the partial fragment of *treY* gene obtained as described above, the DNA primers P16 (SEQ ID NO: 16) and P26 (SEQ ID NO: 26) were newly synthesized, and unknown regions flanking to the partial fragment was amplified by "inverse PCR" (Triglia, T. et al., *Nucleic Acids Res.*, 16, 81-86 (1988); Ochman H., et al., *Genetics*, 120, 621-623 (1988)). The chromosomal DNA of *Brevibacterium lactofermentum* ATCC 13869 was digested with a restriction enzyme *Bam*HI, *Hind*III, *Sal*I or *Xho*I, and self-ligated by using T4 DNA ligase (Takara Shuzo). By using this as a template and the DNA primers P16 and P26, PCR was performed with a cycle consisting of reactions at 94°C for 0.5 minute, 55°C for 1 minute and 72°C for 4 minutes, which was repeated for 30 cycles. As a result, when



*Hind*III or *Sac*I was used as the restriction enzyme, an amplified fragment of 0.6 kbp or 1.5 kbp was obtained, respectively. The nucleotide sequences of these amplified fragments were directly determined by using the DNA primers P16 to P28 (SEQ ID NOS: 16-28). Thus, the entire nucleotide sequence of *treY* gene of *Brevibacterium lactofermentum* ATCC 13869 was determined as shown in SEQ ID NO: 31. The amino acid sequence encoded by this nucleotide sequence is shown in SEQ ID NOS: 31 and 32.

[0054] When homology of the sequence of the aforementioned *treY* gene was determined with respect to the *treY* gene of *Arthrobacter* sp. (GenBank accession D63343), *treY* gene of *Brevibacterium helvolum* (GenBank accession AF039919) and *treY* gene of *Rhizobium* sp. (GenBank accession D78001), the nucleotide sequence showed homologies of 52.0%, 52.3% and 51.9%, respectively, and the amino acid sequence showed homologies of 40.9%, 38.5% and 39.8%, respectively. The homologies were calculated by using software, "GENETIX-WIN" (Software Development), based on the Lipman-Person method (*Science*, 227, 1435-1441 (1985)).

## <2> Preparation of plasmid for *treY* gene disruption

[0055] In order to examine presence or absence of improvement effect in L-glutamic acid productivity by disruption of the gene coding for the enzyme in trehalose biosynthesis pathway in coryneform bacteria, a plasmid for *treY* gene disruption was produced. First, PCR was performed by using the primers P17 (SEQ ID NO: 17) and P25 (SEQ ID NO: 25) and the chromosomal DNA of ATCC 13869 as a template with a cycle consisting of reactions at 94°C for 0.5 minute, 60°C for 0.5 minute and 72°C for 2 minutes, which was repeated for 30 cycles. The amplified fragment was digested with *Eco*RI and ligated to pHSG299 (Takara Shuzo) digested with *Eco*RI by using T4 DNA ligase (Takara Shuzo) to obtain a plasmid pHtreY. Further, this pHtreY was digested with *A*flI (Takara Shuzo), blunt-ended by using T4 DNA polymerase (Takara Shuzo), and self-ligated by using T4 ligase (Takara Shuzo) to construct a plasmid pHtreYA containing the *treY* gene having a frame shift mutation (four nucleotides were inserted after the 1145th nucleotide in the sequence of SEQ ID NO: 31) at an approximately central part thereof.

## <3> Preparation of *treY* gene-disrupted strain

[0056] By using the plasmid pCtreYA for gene disruption, a L-glutamic acid producing bacterium, *Brevibacterium lactofermentum* ATCC 13869, was transformed by the electric pulse method, and transformants were selected as to the ability to grow in CM2B medium containing 20 mg/L of kanamycin. Because the plasmid pCtreYA for *treY* gene disruption does not have a replication origin that could function in *Brevibacterium lactofermentum*, the transformants obtained by using the plasmid suffered recombination occurred between the *treY* genes on the *Brevibacterium lactofermentum* chromosome and the plasmid pCtreYA for gene disruption. From the homologous recombinant strains obtained as described above, strains in which the vector portion of the plasmid pCtreYA for gene disruption was eliminated due to re-occurrence of homologous recombination were selected based on acquired kanamycin sensitivity as a marker.

[0057] From the strains obtained as described above, a strain introduced with the desired frame shift mutation was selected. Selection of such a strain was performed by PCR using the DNA primers P19 (SEQ ID NO: 19) and P25 (SEQ ID NO: 25) with a cycle consisting of reactions at 94°C for 0.5 minute, 55°C for 0.5 minute and 72°C for 1.5 minutes, which was repeated for 30 cycles, and sequencing the obtained fragment using the DNA primer P21 or P23 to confirm dysfunction of the *treY* gene due to introduction of frame shift mutation. The strain obtained as described above was designated as  $\Delta$ TA strain.

## Example 3: Evaluation of L-glutamic acid producing ability of $\Delta$ OA strain and $\Delta$ TA strain

[0058] The ATCC 13869 strain,  $\Delta$ OA strain and  $\Delta$ TA strain were each cultured for producing L-glutamic acid as follows. Each of these strains was refreshed by culturing it on a CM2B plate medium, and each refreshed strain was cultured in a medium containing 80 g of glucose, 1 g of  $\text{KH}_2\text{PO}_4$ , 0.4 g of  $\text{MgSO}_4$ , 30 g of  $(\text{NH}_4)_2\text{SO}_4$ , 0.01 g of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.01 g  $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$ , 15 ml of soybean hydrolysate solution, 200  $\mu\text{g}$  of thiamin hydrochloride, 3  $\mu\text{g}$  of biotin and 50 g of  $\text{CaCO}_3$  in 1 L of pure water (adjusted to pH 8.0 with KOH) at 31.5°C. After the culture, amount of L-glutamic acid accumulated in the medium and absorbance at 620 nm of the culture broth diluted 51 times were measured. The results are shown in Table 1.

[0059] The *Brevibacterium lactofermentum* strains of which *otsA* gene or *treY* gene was disrupted showed growth in a degree similar to that of the parent strain, and in addition, increased L-glutamic acid production compared with the parent strain.

# EP 1 174 508 A2

Table 1

Strain	OD <sub>620</sub> (x51)	L-Glutamic acid (g/L)	Yield (%)
ATCC 13869	0.930	40.2	48.4
ΔOA	1.063	43.8	52.8
ΔTA	0.850	45.6	54.9

(Explanation of sequence Listing)

[0060]

SEQ ID NO: 1: Primer P1 for amplification of *otsA*

SEQ ID NO: 2: Primer P2 for amplification of *otsA*

SEQ ID NO: 3: Primer P5

SEQ ID NO: 4: Primer P6

SEQ ID NO: 5: Primer P7

SEQ ID NO: 6: Primer P8

SEQ ID NO: 7: Primer P9

SEQ ID NO: 8: Primer P10

SEQ ID NO: 9: Primer P11

SEQ ID NO: 10: Primer P12

SEQ ID NO: 11: Primer P13

SEQ ID NO: 12: Primer P14

SEQ ID NO: 13: Primer P15

SEQ ID NO: 14: Primer P3 for amplification of *treY*

SEQ ID NO: 15: Primer P4 for amplification of *treY*

SEQ ID NO: 16: Primer P16

SEQ ID NO: 17: Primer P17

SEQ ID NO: 18: Primer P18

SEQ ID NO: 19: Primer P19

SEQ ID NO: 20: Primer P20

SEQ ID NO: 21: Primer P21

SEQ ID NO: 22: Primer P22

SEQ ID NO: 23: Primer P23

SEQ ID NO: 24: Primer P24

SEQ ID NO: 25: Primer P25

SEQ ID NO: 26: Primer P26

SEQ ID NO: 27: Primer P27

SEQ ID NO: 28: Primer P28

SEQ ID NO: 29: Nucleotide sequence of *otsA* gene

SEQ ID NO: 30: Amino acid sequence of OtsA

SEQ ID NO: 31: Nucleotide sequence of *treY* gene

SEQ ID NO: 32: Amino acid sequence of TreY

SEQ ID NO: 33: Primer P29

SEQ ID NO: 34: Primer P30

SEQUENCE LISTING

5 <110> Ajinomoto Co., Inc.

<120> Bacterium Producing L-Glutamic Acid and Method for Producing L-  
 10 Glutamic Acid

<130> OP1195

15 <140>  
 <141> 2000-07-

20 <150> JP 2000-204256  
 <151> 2000-07-05

25 <160> 34

<170> PatentIn Ver. 2.0

30 <210> 1  
 <211> 20  
 <212> DNA  
 35 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: primer for PCR

40

---

<220>  
 <221> misc\_feature  
 45 <222> (3,9,18)  
 <223> n=a or g or c or t

<400> 1  
 50 canathggnt tyttyytnc

20

<210> 2  
 55

EP 1 174 508 A2

<211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: primer for PCR

<220>  
 <221> misc\_feature  
 <222> (3,11,19)  
 <223> n=a or g or c or t

<400> 2  
 canarrttca tncrctcnc 19

<210> 3  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: primer for PCR

<400> 3  
 gaatcatcca tataagatcc ggc 23

<210> 4  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: primer for PCR

<400> 4  
 tagctttgta gttgttgcta accg 24

<210> 5  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for PCR

5

<400> 5

agcgaacttg aggtttactt cccg

24

10

<210> 6

<211> 24

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: primer for PCR

20

<400> 6

tgctggttcc tggcatttg cgcc

24

25

<210> 7

<211> 20

<212> DNA

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence: primer for PCR

35

<400> 7

tcgaacaatc tcttcacgcc

20

40

<210> 8

<211> 21

<212> DNA

<213> Artificial Sequence

45

<220>

<223> Description of Artificial Sequence: primer for PCR

50

<400> 8

gaatcccacc aaatctgcgc c

21

55

<210> 9

EP 1 174 508 A2

5 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

10 <220>  
 <223> Description of Artificial Sequence: primer for PCR

15 <400> 9  
 tgatgttgaa atgtttgggg 20  
 <210> 10  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

20 <220>  
 <223> Description of Artificial Sequence: primer for PCR  
 <400> 10  
 gatgtcatgc tggttacgcc 20

25 <210> 11  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence

30 <220>  
 <223> Description of Artificial Sequence: primer for PCR

35 <400> 11  
 caaagcacca gtgccgtcgc gg 22

40 <210> 12  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

45 <220>  
 <223> Description of Artificial Sequence: primer for PCR

50 <400> 12  
 tgttcgtttt cattcgcgtt gccg 24

55 <210> 13

EP 1 174 508 A2

5 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

10 <220>  
 <223> Description of Artificial Sequence: primer for PCR

15 <400> 13  
 atagtttcct ggattgttg gcgc 24

20 <210> 14  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

25 <220>  
 <223> Description of Artificial Sequence: primer for PCR

30 <220>  
 <221> misc\_feature  
 <222> (9)  
 <223> n=a or g or c or t

35 <400> 14  
 caraayccnt ggtggtgg 18

40 <210> 15  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

45 <220>  
 <223> Description of Artificial Sequence: primer for PCR

50 <220>  
 <221> misc\_feature  
 <222> (3,6,15)  
 <223> n=a or g or c or t

55 <400> 15  
 ggncgncgrt trtcnggrtc 20



EP 1 174 508 A2

5	<210> 16	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Description of Artificial Sequence: primer for PCR	
15	<400> 16	
	cgagctcttc attgatggcg	20
20	<210> 17	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Description of Artificial Sequence: primer for PCR	
30	<400> 17	
	gcagctacac acgagttggg	20
35	<210> 18	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
40	<220>	
	<223> Description of Artificial Sequence: primer for PCR	
45	<400> 18	
	gcaacaccta aatggttg	20
50	<210> 19	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Description of Artificial Sequence: primer for PCR	

EP 1 174 508 A2

5 <400> 19  
gcaagaagtc tacaagcgcc 20

10 <210> 20  
<211> 16  
<212> DNA  
<213> Artificial Sequence

15 <220>  
<223> Description of Artificial Sequence: primer for PCR

20 <400> 20  
gccaacgtat tcacgg 16

25 <210> 21  
<211> 20  
<212> DNA  
<213> Artificial Sequence

30 <220>  
<223> Description of Artificial Sequence: primer for PCR

35 <400> 21  
tgatgaacca ctcgatcccc 20

40 <210> 22  
<211> 20  
<212> DNA  
<213> Artificial Sequence

45 <220>  
<223> Description of Artificial Sequence: primer for PCR

50 <400> 22  
aagacaccac cttctaccgc 20

55 <210> 23  
<211> 20  
<212> DNA  
<213> Artificial Sequence

	<220>	
5	<223> Description of Artificial Sequence: primer for PCR	
	<400> 23	
10	caagtggaat tctgcagcgg	20
	<210> 24	
	<211> 21	
15	<212> DNA	
	<213> Artificial Sequence	
	<220>	
20	<223> Description of Artificial Sequence: primer for PCR	
	<400> 24	
25	cctcctacaa aacctgctgg g	21
	<210> 25	
	<211> 20	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
35	<223> Description of Artificial Sequence: primer for PCR	
	<400> 25	
40	tcgcggatag cttttagggc	20
	<210> 26	
	<211> 20	
	<212> DNA	
45	<213> Artificial Sequence	
	<220>	
50	<223> Description of Artificial Sequence: primer for PCR	
	<400> 26	
	tgagttttta gaagactccc	20
55	<210> 27	

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for PCR

<400> 27

cgcttcagtg gtgttgtccc

20

<210> 28

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for PCR

<400> 28

cgtaccactc cacggaaatt cccg

24

<210> 29

<211> 2369

<212> DNA

<213> *Brevibacterium lactofermentum*

<220>

<221> CDS

<222> (484)..(1938)

<400> 29

acagaatcag cgccggcaga gaaacgtcca aagactaatc agagattcgg tataaaggta 60  
 aaaatcaacc tgcttaggcg tctttcgctt aaatagcgta gaatatcggg tcgatcgctt 120  
 ttaaacactc aggaggatcc ttgccggcca aaatcacgga cactcgtecc accccagaat 180  
 cccttcacgc tgttgaagag gaaaccgcag ccggtgcccg caggattgtt gccacctatt 240  
 ctaaggactt cttegacggc gtcactttga tgtgcatgct cggcgttgaa cctcagggcc 300  
 tgcgttacac caaggtcgct tctgaacacg aggaagctca gccaaagaag gctacaaagc 360  
 ggactcgtaa ggctaccagc taagaaggct gctgctaaga aaacgaccaa gaagaccact 420  
 aagaaaacta ctaaaaagac caccgcaaag aagaccacaa agaagtctta agccggatct 480  
 tat atg gat gat tcc aat agc ttt gta gtt gtt gct aac cgt ctg cca 528

Met Asp Asp Ser Asn Ser Phe Val Val Val Ala Asn Arg Leu Pro

EP 1 174 508 A2

	1	5	10	15	
	gtg gat atg act gtc cac cca gat ggt agc tat agc atc tcc ccc agc	576			
5	Val Asp Met Thr Val His Pro Asp Gly Ser Tyr Ser Ile Ser Pro Ser				
	20	25	30		
	ccc ggt ggc ctt gtc acg ggg ctt tcc ccc gtt ctg gaa caa cat cgt	624			
	Pro Gly Gly Leu Val Thr Gly Leu Ser Pro Val Leu Glu Gln His Arg				
10	35	40	45		
	gga tgt tgg gtc gga tgg cct gga act gta gat gtt gca ccc gaa cca	672			
	Gly Cys Trp Val Gly Trp Pro Gly Thr Val Asp Val Ala Pro Glu Pro				
	50	55	60		
15	ttt cga aca gat acg ggt gtt ttg ctg cac cct gtt gtc ctc act gca	720			
	Phe Arg Thr Asp Thr Gly Val Leu Leu His Pro Val Val Leu Thr Ala				
	65	70	75		
20	agt gac tat gaa ggc ttc tac gag ggc ttt tca aac gca acg ctg tgg	768			
	Ser Asp Tyr Glu Gly Phe Tyr Glu Gly Phe Ser Asn Ala Thr Leu Trp				
	80	85	90	95	
	cct ctt ttc cac gat ctg att gtt act ccg gtg tac aac acc gat tgg	816			
25	Pro Leu Phe His Asp Leu Ile Val Thr Pro Val Tyr Asn Thr Asp Trp				
	100	105	110		
	tgg cat gcg ttt cgg gaa gta aac ctc aag ttc gct gaa gcc gtg agc	864			
	Trp His Ala Phe Arg Glu Val Asn Leu Lys Phe Ala Glu Ala Val Ser				
30	115	120	125		
	caa gtg gcg gca cac ggt gcc act gtg tgg gtg cag gac tat cag ctg	912			
	Gln Val Ala Ala His Gly Ala Thr Val Trp Val Gln Asp Tyr Gln Leu				
	130	135	140		
35	ttg ctg gtt cct ggc att ttg cgc cag atg cgc ctt gat ttg aag atc	960			
	Leu Leu Val Pro Gly Ile Leu Arg Gln Met Arg Leu Asp Leu Lys Ile				
	145	150	155		
	ggt ttc ttc ctc cac att ccc ttc cct tcc cct gat ctg ttc cgt cag	1008			
40	Gly Phe Phe Leu His Ile Pro Phe Pro Ser Pro Asp Leu Phe Arg Gln				
	<del>160</del>	<del>165</del>	<del>170</del>	<del>175</del>	
	ctg ccg tgg cgt gaa gag att gtt cga ggc atg ctg ggc gca gat ttg	1056			
45	Leu Pro Trp Arg Glu Glu Ile Val Arg Gly Met Leu Gly Ala Asp Leu				
	180	185	190		
	gtg gga ttc cat ttg gtt caa aac gca gaa aac ttc ctt gcg tta acc	1104			
	Val Gly Phe His Leu Val Gln Asn Ala Glu Asn Phe Leu Ala Leu Thr				
	195	200	205		
50	cag cag gtt gcc ggc act gcc ggg tct cat gtg ggt cag ccg gac acc	1152			
	Gln Gln Val Ala Gly Thr Ala Gly Ser His Val Gly Gln Pro Asp Thr				
	210	215	220		
55	ttg cag gtc agt ggt gaa gca ttg gtg cgt gag att ggc gct cat gtt	1200			

EP 1 174 508 A2

	Leu Gln Val Ser Gly Glu Ala Leu Val Arg Glu Ile Gly Ala His Val	
	225 230 235	
5	gaa acc gct gac gga agg cga gtt agc gtc ggg gcg ttc ccg atc tcg	1248
	Glu Thr Ala Asp Gly Arg Arg Val Ser Val Gly Ala Phe Pro Ile Ser	
	240 245 250 255	
10	att gat gtt gaa atg ttt ggg gag gcg tcg aaa agc gcc gtt ctt gat	1296
	Ile Asp Val Glu Met Phe Gly Glu Ala Ser Lys Ser Ala Val Leu Asp	
	260 265 270	
	ctt tta aaa acg ctc gac gag ccg gaa acc gta ttc ctg ggc gtt gac	1344
15	Leu Leu Lys Thr Leu Asp Glu Pro Glu Thr Val Phe Leu Gly Val Asp	
	275 280 285	
	cga ctg gac tac acc aag ggc att ttg cag cgc ctg ctt gcg ttt gag	1392
	Arg Leu Asp Tyr Thr Lys Gly Ile Leu Gln Arg Leu Leu Ala Phe Glu	
20	290 295 300	
	gaa ctg ctg gaa tcc ggc gcg ttg gag gcc gac aaa gct gtg ttg ctg	1440
	Glu Leu Leu Glu Ser Gly Ala Leu Glu Ala Asp Lys Ala Val Leu Leu	
	305 310 315	
25	cag gtc gcg acg cct tcg cgt gag cgc att gat cac tat cgt gtg tcg	1488
	Gln Val Ala Thr Pro Ser Arg Glu Arg Ile Asp His Tyr Arg Val Ser	
	320 325 330 335	
	cgt tcg cag gtc gag gaa gcc gtc ggc cgt atc aat ggt cgt ttc ggt	1536
30	Arg Ser Gln Val Glu Glu Ala Val Gly Arg Ile Asn Gly Arg Phe Gly	
	340 345 350	
	cgc atg ggg cgt ccc gtg gtg cat tat cta cac agg tca ttg agc aaa	1584
	Arg Met Gly Arg Pro Val Val His Tyr Leu His Arg Ser Leu Ser Lys	
35	355 360 365	
	aat gat ctc cag gtg ctg tat acc gca gcc gat gtc atg ctg gtt acg	1632
	Asn Asp Leu Gln Val Leu Tyr Thr Ala Ala Asp Val Met Leu Val Thr	
	370 375 380	
40	cct ttt aaa gac ggt atg aac ttg gtg gct aaa gaa ttc gtg gcc aac	1680
	Pro Phe Lys Asp Gly Met Asn Leu Val Ala Lys Glu Phe Val Ala Asn	
	385 390 395	
45	cac cgc gac ggc act ggt gct ttg gtg ctg tcc gaa ttt gcc ggc gcg	1728
	His Arg Asp Gly Thr Gly Ala Leu Val Leu Ser Glu Phe Ala Gly Ala	
	400 405 410 415	
	gcc act gag ctg acc ggt gcg tat tta tgc aac cca ttt gat gtg gaa	1776
50	Ala Thr Glu Leu Thr Gly Ala Tyr Leu Cys Asn Pro Phe Asp Val Glu	
	420 425 430	
	tcc atc aaa cgg caa atg gtg gca gct gtc cat gat ttg aag cac aat	1824
	Ser Ile Lys Arg Gln Met Val Ala Ala Val His Asp Leu Lys His Asn	
55	435 440 445	

EP 1 174 508 A2

ccg gaa tct gcg gca acg cga atg aaa acg aac agc gag cag gtc tat 1872  
 Pro Glu Ser Ala Ala Thr Arg Met Lys Thr Asn Ser Glu Gln Val Tyr  
 5                   450                   455                   460  
 acc cac gac gtc aac gtg tgg gct aat agt ttc ctg gat tgt ttg gcg 1920  
 Thr His Asp Val Asn Val Trp Ala Asn Ser Phe Leu Asp Cys Leu Ala  
                  465                   470                   475  
 10   cag tcg gga gaa aac tca tgaaccgcgc acgaatcgcg accataggcg 1968  
      Gln Ser Gly Glu Asn Ser  
      480                   485  
 15   ttcttcgct tgctttactg ctggcgctct gtggttcaga caccgtggaa atgacagatt 2028  
      ccacctggtt ggtgaccaat atttacaccg atccagatga gtcgaattcg atcagtaatc 2088  
      ttgtcatttc ccagcccagc ttagattttg gcaattcttc cctgtctggt ttactggct 2148  
      gtgtgccttt tacggggcgt gcggaattct tccaaatgg tgagcaaagc tctgttctgg 2208  
      atgccgatta tgtgacctg tcttcctgg atttcgataa acttcocgat gattgccaag 2268  
 20   gacaagaact caaagttcat aacgagctgg ttgatcttct gcctggttct ttgaaatct 2328  
      ccaggacttc tggttcagaa atcttgctga ctacgatgt c 2369  
  
 25   <210> 30  
      <211> 485  
      <212> PRT  
      <213> Brevibacterium lactofermentum  
  
 30   <400> 30  
      Met Asp Asp Ser Asn Ser Phe Val Val Val Ala Asn Arg Leu Pro Val  
      1                   5                   10                   15  
 35   Asp Met Thr Val His Pro Asp Gly Ser Tyr Ser Ile Ser Pro Ser Pro  
                  20                   25                   30  
      Gly Gly Leu Val Thr Gly Leu Ser Pro Val Leu Glu Gln His Arg Gly  
                  35                   40                   45  
 40   Cys Trp Val Gly Trp Pro Gly Thr Val Asp Val Ala Pro Glu Pro Phe  
      50                   55                   60  
      Arg Thr Asp Thr Gly Val Leu Leu His Pro Val Val Leu Thr Ala Ser  
      65                   70                   75                   80  
 45   Asp Tyr Glu Gly Phe Tyr Glu Gly Phe Ser Asn Ala Thr Leu Trp Pro  
                  85                   90                   95  
      Leu Phe His Asp Leu Ile Val Thr Pro Val Tyr Asn Thr Asp Trp Trp  
                  100                   105                   110  
 50   His Ala Phe Arg Glu Val Asn Leu Lys Phe Ala Glu Ala Val Ser Gln  
                  115                   120                   125  
      Val Ala Ala His Gly Ala Thr Val Trp Val Gln Asp Tyr Gln Leu Leu  
      130                   135                   140  
 55



EP 1 174 508 A2

5	Leu Val Pro Gly Ile Leu Arg Gln Met Arg Leu Asp Leu Lys Ile Gly	145	150	155	160
	Phe Phe Leu His Ile Pro Phe Pro Ser Pro Asp Leu Phe Arg Gln Leu	165	170	175	
10	Pro Trp Arg Glu Glu Ile Val Arg Gly Met Leu Gly Ala Asp Leu Val	180	185	190	
	Gly Phe His Leu Val Gln Asn Ala Glu Asn Phe Leu Ala Leu Thr Gln	195	200	205	
15	Gln Val Ala Gly Thr Ala Gly Ser His Val Gly Gln Pro Asp Thr Leu	210	215	220	
	Gln Val Ser Gly Glu Ala Leu Val Arg Glu Ile Gly Ala His Val Glu	225	230	235	240
20	Thr Ala Asp Gly Arg Arg Val Ser Val Gly Ala Phe Pro Ile Ser Ile	245	250	255	
	Asp Val Glu Met Phe Gly Glu Ala Ser Lys Ser Ala Val Leu Asp Leu	260	265	270	
25	Leu Lys Thr Leu Asp Glu Pro Glu Thr Val Phe Leu Gly Val Asp Arg	275	280	285	
	Leu Asp Tyr Thr Lys Gly Ile Leu Gln Arg Leu Leu Ala Phe Glu Glu	290	295	300	
30	Leu Leu Glu Ser Gly Ala Leu Glu Ala Asp Lys Ala Val Leu Leu Gln	305	310	315	320
	Val Ala Thr Pro Ser Arg Glu Arg Ile Asp His Tyr Arg Val Ser Arg	325	330	335	
35	Ser Gln Val Glu Glu Ala Val Gly Arg Ile Asn Gly Arg Phe Gly Arg	340	345	350	
	Met Gly Arg Pro Val Val His Tyr Leu His Arg Ser Leu Ser Lys Asn	355	360	365	
40	Asp Leu Gln Val Leu Tyr Thr Ala Ala Asp Val Met Leu Val Thr Pro	370	375	380	
45	Phe Lys Asp Gly Met Asn Leu Val Ala Lys Glu Phe Val Ala Asn His	385	390	395	400
	Arg Asp Gly Thr Gly Ala Leu Val Leu Ser Glu Phe Ala Gly Ala Ala	405	410	415	
50	Thr Glu Leu Thr Gly Ala Tyr Leu Cys Asn Pro Phe Asp Val Glu Ser	420	425	430	
	Ile Lys Arg Gln Met Val Ala Ala Val His Asp Leu Lys His Asn Pro	435	440	445	
55	Glu Ser Ala Ala Thr Arg Met Lys Thr Asn Ser Glu Gln Val Tyr Thr	450	455	460	
	His Asp Val Asn Val Trp Ala Asn Ser Phe Leu Asp Cys Leu Ala Gln				

EP 1 174 508 A2

465                      470                      475                      480  
Ser Gly Glu Asn Ser  
5                                      485

<210> 31  
<211> 2956  
10 <212> DNA  
<213> Brevibacterium lactofermentum

<220>  
15 <221> CDS  
<222> (82)..(2514)

<220>  
20 <221> misc\_feature  
<222> (2953)  
<223> n=a or g or c or t

25 <400> 31  
ttttccacag cagggaaggc gtgaacacta agatcgagga cgtaccgcac gattttgcct 60  
aacttttaag ggtgtttcat c atg gca cgt cca att tcc gca acg tac agg 111  
30                                      Met Ala Arg Pro Ile Ser Ala Thr Tyr Arg  
   1                                      5                                      10  
ctt caa atg cga gga cct caa gca gat agc gcc ggg cgt ttc ttt ggt 159  
Leu Gln Met Arg Gly Pro Gln Ala Asp Ser Ala Gly Arg Phe Phe Gly  
35                                      15                                      20                                      25  
ttt gcg cag gcc aaa gcc cag ctt ccc tat ctg aag aag cta ggc atc 207  
Phe Ala Gln Ala Lys Ala Gln Leu Pro Tyr Leu Lys Lys Leu Gly Ile  
   30                                      35                                      40  
40 agc cac ctg tac ctc tcc cct att ttt acg gcc atg cca gat tcc aat 255  
Ser His Leu Tyr Leu Ser Pro Ile Phe Thr Ala Met Pro Asp Ser Asn  
   45                                      50                                      55  
cat ggc tac gat gtc att gat ccc acc gcc atc aat gaa gag ctc ggt 303  
45 His Gly Tyr Asp Val Ile Asp Pro Thr Ala Ile Asn Glu Glu Leu Gly  
   60                                      65                                      70  
ggc atg gag ggt ctt cga gat ctt gct gca gct aca cac gag ttg ggc 351  
Gly Met Glu Gly Leu Arg Asp Leu Ala Ala Ala Thr His Glu Leu Gly  
50                                      75                                      80                                      85                                      90  
atg ggc atc atc att gat att gtt ccc aac cat tta ggt gtt gcc gtt 399  
Met Gly Ile Ile Ile Asp Ile Val Pro Asn His Leu Gly Val Ala Val  
55                                      95                                      100                                      105

EP 1 174 508 A2

	cca cat ttg aat cct tgg tgg tgg gat gtt cta aaa aac ggc aaa gat	447
	Pro His Leu Asn Pro Trp Trp Trp Asp Val Leu Lys Asn Gly Lys Asp	
5	110 115 120	
	tcc gct ttt gag ttc tat ttc gat att gac tgg cac gaa gac aac ggt	495
	Ser Ala Phe Glu Phe Tyr Phe Asp Ile Asp Trp His Glu Asp Asn Gly	
	125 130 135	
10	tct ggt ggc aag ctg ggc atg ccg att ctg ggt gct gaa ggc gat gaa	543
	Ser Gly Gly Lys Leu Gly Met Pro Ile Leu Gly Ala Glu Gly Asp Glu	
	140 145 150	
15	gac aag ctg gaa ttc gcg gag ctt gat gga gag aaa gtg ctc aaa tat	591
	Asp Lys Leu Glu Phe Ala Glu Leu Asp Gly Glu Lys Val Leu Lys Tyr	
	155 160 165 170	
	ttt gac cac ctc ttc cca atc gcg cct ggt acc gaa gaa ggg aca ccg	639
	Phe Asp His Leu Phe Pro Ile Ala Pro Gly Thr Glu Glu Gly Thr Pro	
20	175 180 185	
	caa gaa gtc tac aag cgc cag cat tac cgc ctg cag ttc tgg cgc gac	687
	Gln Glu Val Tyr Lys Arg Gln His Tyr Arg Leu Gln Phe Trp Arg Asp	
	190 195 200	
25	ggc gtg atc aac ttc cgt cgc ttc ttt tcc gtg aat acg ttg gct ggc	735
	Gly Val Ile Asn Phe Arg Arg Phe Phe Ser Val Asn Thr Leu Ala Gly	
	205 210 215	
30	atc agg caa gaa gat ccc ttg gtg ttt gaa cat act cat cgt ctg ctg	783
	Ile Arg Gln Glu Asp Pro Leu Val Phe Glu His Thr His Arg Leu Leu	
	220 225 230	
	cgc gaa ttg gtg gcg gaa gac ctc att gac ggc gtg cgc gtc gat cac	831
	Arg Glu Leu Val Ala Glu Asp Leu Ile Asp Gly Val Arg Val Asp His	
35	235 240 245 250	
	ccc gac ggg ctt tcc gat cct ttt gga tat ctg cac aga ctc cgc gac	879
	Pro Asp Gly Leu Ser Asp Pro Phe Gly Tyr Leu His Arg Leu Arg Asp	
	255 260 265	
40	ctc att gga cct gac cgc tgg ctg atc atc gaa aag atc ttg agc gtt	927
	Leu Ile Gly Pro Asp Arg Trp Leu Ile Ile Glu Lys Ile Leu Ser Val	
	270 275 280	
45	gat gaa cca ctc gat ccc cgc ctg gcc gtt gat ggc acc act ggc tac	975
	Asp Glu Pro Leu Asp Pro Arg Leu Ala Val Asp Gly Thr Thr Gly Tyr	
	285 290 295	
	gac ccc ctc cgt gaa ctc gac ggc gtg ttt atc tcc cga gaa tct gag	1023
	Asp Pro Leu Arg Glu Leu Asp Gly Val Phe Ile Ser Arg Glu Ser Glu	
50	300 305 310	
	gac aaa ttc tcc atg ttg gcg ctg acc cac agt gga tcc acc tgg gat	1071
	Asp Lys Phe Ser Met Leu Ala Leu Thr His Ser Gly Ser Thr Trp Asp	

55

EP 1 174 508 A2

	315	320	325	330	
	gaa cgc gcc cta aaa tcc acg gag gaa agc ctc aaa cga gtc gtc gcg				1119
5	Glu Arg Ala Leu Lys Ser Thr Glu Glu Ser Leu Lys Arg Val Val Ala				
	335	340	345		
	caa caa gaa ctc gca gcc gaa atc tta agg ctc gcc cgc gcc atg cgc				1167
10	Gln Gln Glu Leu Ala Ala Glu Ile Leu Arg Leu Ala Arg Ala Met Arg				
	350	355	360		
	cgc gat aac ttc tcc acc gca ggc acc aac gtc acc gaa gac aaa ctt				1215
	Arg Asp Asn Phe Ser Thr Ala Gly Thr Asn Val Thr Glu Asp Lys Leu				
15	365	370	375		
	agc gaa acc atc atc gaa tta gtc gcc gcc atg ccc gtc tac cgc gcc				1263
	Ser Glu Thr Ile Ile Glu Leu Val Ala Ala Met Pro Val Tyr Arg Ala				
	380	385	390		
20	gac tac atc tcc ctc tca cgc acc acc gcc acc gtc atc gcg gag atg				1311
	Asp Tyr Ile Ser Leu Ser Arg Thr Thr Ala Thr Val Ile Ala Glu Met				
	395	400	405	410	
	tcc aaa cgc ttc ccc tcc cgg cgc gac gca ctc gac ctc atc tgc gcc				1359
25	Ser Lys Arg Phe Pro Ser Arg Arg Asp Ala Leu Asp Leu Ile Ser Ala				
	415	420	425		
	gcc cta ctt ggc aat ggc gag gcc aaa atc cgc ttc gcc caa gtc tgc				1407
	Ala Leu Leu Gly Asn Gly Glu Ala Lys Ile Arg Phe Ala Gln Val Cys				
30	430	435	440		
	ggc gcc gtc atg gcc aaa ggt gtg gaa gac acc acc ttc tac cgc gca				1455
	Gly Ala Val Met Ala Lys Gly Val Glu Asp Thr Thr Phe Tyr Arg Ala				
	445	450	455		
35	tct agg ctc gtt gca ctg caa gaa gtc ggt ggc gcg ccg ggc agg ttc				1503
	Ser Arg Leu Val Ala Leu Gln Glu Val Gly Gly Ala Pro Gly Arg Phe				
	460	465	470		
40	ggc gtc tcc got gca gaa ttc cac ttg ctg cag gaa gaa cgc agc ctg				1551
	Gly Val Ser Ala Ala Glu Phe His Leu Leu Gln Glu Glu Arg Ser Leu				
	475	480	485	490	
	ctg tgg cca cgc acc atg acc acc ttg tcc acg cac gac acc aaa cgc				1599
45	Leu Trp Pro Arg Thr Met Thr Thr Leu Ser Thr His Asp Thr Lys Arg				
	495	500	505		
	ggc gaa gat acc cgc gcc cgc atc atc tcc ctg tcc gaa gtc ccc gat				1647
	Gly Glu Asp Thr Arg Ala Arg Ile Ile Ser Leu Ser Glu Val Pro Asp				
50	510	515	520		
	atg tac tcc gag ctg gtc aat cgt gtt ttc gca gtg ctc ccc gcg cca				1695
	Met Tyr Ser Glu Leu Val Asn Arg Val Phe Ala Val Leu Pro Ala Pro				
	525	530	535		
55	gac ggc gca acg ggc agt ttc ctc cta caa aac ctg ctg ggc gta tgg				1743

EP 1 174 508 A2

	Asp	Gly	Ala	Thr	Gly	Ser	Phe	Leu	Leu	Gln	Asn	Leu	Leu	Gly	Val	Trp	
	540						545					550					
5	ccc	gcc	gac	ggc	gtg	atc	acc	gat	gcg	ctg	cgc	gat	cga	ttc	agg	gaa	1791
	Pro	Ala	Asp	Gly	Val	Ile	Thr	Asp	Ala	Leu	Arg	Asp	Arg	Phe	Arg	Glu	
	555						560					565				570	
	tac	gcc	cta	aaa	gct	atc	cgc	gaa	gca	tcc	aca	aaa	acc	acg	tgg	gtg	1839
10	Tyr	Ala	Leu	Lys	Ala	Ile	Arg	Glu	Ala	Ser	Thr	Lys	Thr	Thr	Trp	Val	
					575					580					585		
	gac	ccc	aac	gag	tcc	ttc	gag	gct	gcg	gtc	tgc	gat	tgg	gtg	gaa	gcg	1887
	Asp	Pro	Asn	Glu	Ser	Phe	Glu	Ala	Ala	Val	Cys	Asp	Trp	Val	Glu	Ala	
15				590					595					600			
	ctt	ttc	gac	gga	ccc	tcc	acc	tca	tta	atc	acc	gaa	ttt	gtc	tcc	cac	1935
	Leu	Phe	Asp	Gly	Pro	Ser	Thr	Ser	Leu	Ile	Thr	Glu	Phe	Val	Ser	His	
		605						610					615				
20	atc	aac	cgt	ggc	tct	gtg	aat	atc	tcc	tta	ggc	agg	aaa	ctg	ctg	caa	1983
	Ile	Asn	Arg	Gly	Ser	Val	Asn	Ile	Ser	Leu	Gly	Arg	Lys	Leu	Leu	Gln	
		620					625					630					
	atg	gtg	ggc	gct	gga	atc	ccc	gac	act	tac	caa	gga	act	gag	ttt	tta	2031
25	Met	Val	Gly	Ala	Gly	Ile	Pro	Asp	Thr	Tyr	Gln	Gly	Thr	Glu	Phe	Leu	
		635				640					645				650		
	gaa	gac	tcc	ctg	gta	gat	ccc	gat	aac	cga	cgc	ttt	gtt	gat	tac	acc	2079
	Glu	Asp	Ser	Leu	Val	Asp	Pro	Asp	Asn	Arg	Arg	Phe	Val	Asp	Tyr	Thr	
30				655						660					665		
	gcc	aga	gaa	caa	gtc	ctg	gag	cgc	ctg	caa	acc	tgg	gat	tgg	acg	cag	2127
	Ala	Arg	Glu	Gln	Val	Leu	Glu	Arg	Leu	Gln	Thr	Trp	Asp	Trp	Thr	Gln	
				670					675					680			
35	gtt	aat	tcg	gta	gaa	gac	ttg	gtg	gat	aac	gcc	gac	atc	gcc	aaa	atg	2175
	Val	Asn	Ser	Val	Glu	Asp	Leu	Val	Asp	Asn	Ala	Asp	Ile	Ala	Lys	Met	
		685					690					695					
	gcc	gtg	gtc	cat	aaa	tcc	ctc	gag	ttg	cgt	gct	gaa	ttt	cgt	gca	agc	2223
40	Ala	Val	Val	His	Lys	Ser	Leu	Glu	Leu	Arg	Ala	Glu	Phe	Arg	Ala	Ser	
	700						705					710					
	ttt	gtt	ggt	gga	gat	cat	cag	gca	gta	ttt	ggc	gaa	ggt	cgc	gca	gaa	2271
45	Phe	Val	Gly	Gly	Asp	His	Gln	Ala	Val	Phe	Gly	Glu	Gly	Arg	Ala	Glu	
	715					720						725			730		
	tcc	cac	atc	atg	ggc	atc	gcc	cgc	ggt	aca	gac	cga	aac	cac	ctc	aac	2319
	Ser	His	Ile	Met	Gly	Ile	Ala	Arg	Gly	Thr	Asp	Arg	Asn	His	Le		

EP 1 174 508 A2

gga tgg tat gac acc acc gtc acg ctt cct ggt gga caa tgg gaa gac 2415  
 Gly Trp Tyr Asp Thr Thr Val Thr Leu Pro Gly Gly Gln Trp Glu Asp  
 5 765 770 775

agg ctc acc ggg caa cgc ttc agt ggt gtt gtc cca gcc acc gat ttg 2463  
 Arg Leu Thr Gly Gln Arg Phe Ser Gly Val Val Pro Ala Thr Asp Leu  
 780 785 790

10 ttc tca cat tta ccc gta tct ttg ttg gtt tta gta ccc gat agt gag 2511  
 Phe Ser His Leu Pro Val Ser Leu Leu Val Leu Val Pro Asp Ser Glu  
 795 800 805 810

15 ttt tgatccctgc acaggaaagt tagcggcgct actatgaacg atcgatatgt 2564  
 Phe

ctgacaacac tctctcccaa tttggcagtt actaccacga attccgacgt gcccatccca 2624  
 tggccgacgt cgaattcttc ctagcaattg aagaattact cacagacggt ggtgtcacct 2684  
 20 tcgatcgcgt caccacacgc atcaaagaat ggtcaagcct gaaagccaag gctcgcgaag 2744  
 gtcgcgacga tggtcgttg atctaccctg atccgcgcaa agacatccac gacatgatcg 2804  
 gtgttcggat caccacgtac cactccacgg aaattcccgt ggccttaaaa gtgtccaag 2864  
 actccttcat cgtccacaaa tccgtagaca aagccgctga aactcgcac tcaggcggt 2924  
 25 ttggttacgg ctcccaccac caaggattnt ag 2956

<210> 32  
 <211> 811  
 30 <212> PRT  
 <213> Brevibacterium lactofermentum

<400> 32

35 Met Ala Arg Pro Ile Ser Ala Thr Tyr Arg Leu Gln Met Arg Gly Pro  
 1 5 10 15  
 Gln Ala Asp Ser Ala Gly Arg Phe Phe Gly Phe Ala Gln Ala Lys Ala  
 20 25 30

40 Gln Leu Pro Tyr Leu Lys Lys Leu Gly Ile Ser His Leu Tyr Leu Ser  
 35 40 45

Pro Ile Phe Thr Ala Met Pro Asp Ser Asn His Gly Tyr Asp Val Ile  
 50 55 60

45 Asp Pro Thr Ala Ile Asn Glu Glu Leu Gly Gly Met Glu Gly Leu Arg  
 65 70 75 80

Asp Leu Ala Ala Ala Thr His Glu Leu Gly Met Gly Ile Ile Ile Asp  
 85 90 95

50 Ile Val Pro Asn His Leu Gly Val Ala Val Pro His Leu Asn Pro Trp  
 100 105 110

Trp Trp Asp Val Leu Lys Asn Gly Lys Asp Ser Ala Phe Glu Phe Tyr  
 115 120 125

55

5	Phe Asp Ile Asp Trp His Glu Asp Asn Gly Ser Gly Gly Lys Leu Gly	130	135	140
	Met Pro Ile Leu Gly Ala Glu Gly Asp Glu Asp Lys Leu Glu Phe Ala	145	150	155
10	Glu Leu Asp Gly Glu Lys Val Leu Lys Tyr Phe Asp His Leu Phe Pro	165	170	175
	Ile Ala Pro Gly Thr Glu Glu Gly Thr Pro Gln Glu Val Tyr Lys Arg	180	185	190
15	Gln His Tyr Arg Leu Gln Phe Trp Arg Asp Gly Val Ile Asn Phe Arg	195	200	205
	Arg Phe Phe Ser Val Asn Thr Leu Ala Gly Ile Arg Gln Glu Asp Pro	210	215	220
20	Leu Val Phe Glu His Thr His Arg Leu Leu Arg Glu Leu Val Ala Glu	225	230	235
	Asp Leu Ile Asp Gly Val Arg Val Asp His Pro Asp Gly Leu Ser Asp	245	250	255
25	Pro Phe Gly Tyr Leu His Arg Leu Arg Asp Leu Ile Gly Pro Asp Arg	260	265	270
	Trp Leu Ile Ile Glu Lys Ile Leu Ser Val Asp Glu Pro Leu Asp Pro	275	280	285
30	Arg Leu Ala Val Asp Gly Thr Thr Gly Tyr Asp Pro Leu Arg Glu Leu	290	295	300
	Asp Gly Val Phe Ile Ser Arg Glu Ser Glu Asp Lys Phe Ser Met Leu	305	310	315
35	Ala Leu Thr His Ser Gly Ser Thr Trp Asp Glu Arg Ala Leu Lys Ser	325	330	335
	Thr Glu Glu Ser Leu Lys Arg Val Val Ala Gln Gln Glu Leu Ala Ala	340	345	350
40	Glu Ile Leu Arg Leu Ala Arg Ala Met Arg Arg Asp Asn Phe Ser Thr	355	360	365
	Ala Gly Thr Asn Val Thr Glu Asp Lys Leu Ser Glu Thr Ile Ile Glu	370	375	380
45	Leu Val Ala Ala Met Pro Val Tyr Arg Ala Asp Tyr Ile Ser Leu Ser	385	390	395
	Arg Thr Thr Ala Thr Val Ile Ala Glu Met Ser Lys Arg Phe Pro Ser	405	410	415
50	Arg Arg Asp Ala Leu Asp Leu Ile Ser Ala Ala Leu Leu Gly Asn Gly	420	425	430
	Glu Ala Lys Ile Arg Phe Ala Gln Val Cys Gly Ala Val Met Ala Lys	435	440	445
55	Gly Val Glu Asp Thr Thr Phe Tyr Arg Ala Ser Arg Leu Val Ala Leu			



EP 1 174 508 A2

	450		455		460	
5	Gln Glu Val Gly Gly Ala Pro Gly Arg Phe Gly Val Ser Ala Ala Glu					
	465		470		475	480
	Phe His Leu Leu Gln Glu Glu Arg Ser Leu Leu Trp Pro Arg Thr Met					
		485		490		495
10	Thr Thr Leu Ser Thr His Asp Thr Lys Arg Gly Glu Asp Thr Arg Ala					
		500		505		510
	Arg Ile Ile Ser Leu Ser Glu Val Pro Asp Met Tyr Ser Glu Leu Val					
		515		520		525
15	Asn Arg Val Phe Ala Val Leu Pro Ala Pro Asp Gly Ala Thr Gly Ser					
		530		535		540
	Phe Leu Leu Gln Asn Leu Leu Gly Val Trp Pro Ala Asp Gly Val Ile					
		545		550		555
20	Thr Asp Ala Leu Arg Asp Arg Phe Arg Glu Tyr Ala Leu Lys Ala Ile					
		565		570		575
	Arg Glu Ala Ser Thr Lys Thr Thr Trp Val Asp Pro Asn Glu Ser Phe					
		580		585		590
25	Glu Ala Ala Val Cys Asp Trp Val Glu Ala Leu Phe Asp Gly Pro Ser					
		595		600		605
	Thr Ser Leu Ile Thr Glu Phe Val Ser His Ile Asn Arg Gly Ser Val					
30		610		615		620
	Asn Ile Ser Leu Gly Arg Lys Leu Leu Gln Met Val Gly Ala Gly Ile					
		625		630		635
	Pro Asp Thr Tyr Gln Gly Thr Glu Phe Leu Glu Asp Ser Leu Val Asp					
		645		650		655
35	Pro Asp Asn Arg Arg Phe Val Asp Tyr Thr Ala Arg Glu Gln Val Leu					
		660		665		670
	Glu Arg Leu Gln Thr Trp Asp Trp Thr Gln Val Asn Ser Val Glu Asp					
40		675		680		685
	Leu Val Asp Asn Ala Asp Ile Ala Lys Met Ala Val Val His Lys Ser					
		690		695		700
45	Leu Glu Leu Arg Ala Glu Phe Arg Ala Ser Phe Val Gly Gly Asp His					
		705		710		715
	Gln Ala Val Phe Gly Glu Gly Arg Ala Glu Ser His Ile Met Gly Ile					
		725		730		735
50	Ala Arg Gly Thr Asp Arg Asn His Leu Asn Ile Ile Ala Leu Ala Thr					
		740		745		750
	Arg Arg Pro Leu Ile Leu Glu Asp Arg Gly Gly Trp Tyr Asp Thr Thr					
		755		760		765
55	Val Thr Leu Pro Gly Gly Gln Trp Glu Asp Arg Leu Thr Gly Gln Arg					
		770		775		780

Phe Ser Gly Val Val Pro Ala Thr Asp Leu Phe Ser His Leu Pro Val  
 785                      790                      795                      800  
 Ser Leu Leu Val Leu Val Pro Asp Ser Glu Phe  
                     805                      810

<210> 33  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: primer for PCR

<400> 33  
 ccaaaatcga taacatcaat cgagatcggg                      30

<210> 34  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: primer for PCR

<400> 34  
 ctgatcgat taaaaacgct cgacgagccg                      30

# Claims

1. A coryneform bacterium having L-glutamic acid producing ability, wherein trehalose synthesis ability is decreased or deleted in the bacterium.
2. The coryneform bacteria according to claim 1, wherein the trehalose synthesis ability is decreased or deleted by introducing a mutation into a chromosomal gene coding for an enzyme in trehalose synthesis pathway or disrupting the gene.
3. The coryneform bacteria according to claim 2, wherein the gene coding for the enzyme in trehalose synthesis pathway consists of a gene coding for trehalose-6-phosphate synthase, a gene coding for maltotriose synthase, or both of these genes.
4. The coryneform bacteria according to claim 3, wherein the gene coding for trehalose-6-phosphate synthase codes for the amino acid sequence of SEQ ID NO: 30, and the gene coding for maltotriose synthase codes

for the amino acid sequence of SEQ ID NO: 32.

5. A method for producing L-glutamic acid comprising the steps of culturing a coryneform bacterium according to any one of claims 1-4 in a medium to produce and accumulate L-glutamic acid in the medium, and collecting the L-glutamic acid from the medium.

6. A DNA coding for a protein defined in the following (A) or (B):

(A) a protein having the amino acid sequence of SEQ ID NO: 30,  
(B) a protein having the amino acid sequence of SEQ ID NO: 30 including substitution, deletion, insertion or addition of one or several amino acid residues and having trehalose-6-phosphate synthase activity.

7. A DNA according to claim 6, which is a DNA defined in the following (a) or (b):

(a) a DNA containing a nucleotide sequence comprising at least the residues of nucleotide numbers 484-1938 in the nucleotide sequence of SEQ ID NO: 29,  
(b) a DNA hybridizable with a nucleotide sequence comprising at least the residues of nucleotide numbers 484-1938 in the nucleotide sequence of SEQ ID NO: 29 under a stringent condition, showing homology of 55% or more to the foregoing nucleotide sequence, and coding for a protein having trehalose-6-phosphate synthase activity.

8. A DNA coding for a protein defined in the following (A) or (B):

(A) a protein having the amino acid sequence of SEQ ID NO: 32,  
(B) a protein having the amino acid sequence of SEQ ID NO: 32 including substitution, deletion, insertion or addition of one or several amino acid residues and having maltotigosyltrehalose synthase activity.

9. A DNA according to claim 8, which is a DNA defined in the following (a) or (b):

(a) a DNA containing a nucleotide sequence comprising at least the residues of nucleotide numbers 82-2514 in the nucleotide sequence of SEQ ID NO: 31,  
(b) a DNA hybridizable with a nucleotide sequence comprising at least the residues of nucleotide numbers 82-2514 in the nucleotide sequence of SEQ ID NO: 31 under a stringent condition, showing homology of 60% or more to the foregoing nucleotide sequence, and coding for a protein having maltotigosyltrehalose synthase activity.